

Reply to “Comment on ‘Properties of the massive Thirring model from the XYZ spin chain’ ”

M. Kolanović*

*Center for Cosmology and Particle Physics, Department of Physics
New York University, New York, NY 10003*

S. Pallua† and P. Prester‡

*Department of Theoretical Physics, PMF, University of Zagreb
Bijenička c. 32, p.p. 331, 10000 Zagreb, Croatia*

(Dated: February 12, 2003)

We elaborate in more details why lattice calculation in [Kolanovic et al, Phys. Rev. D **62**, 025021 (2000)] was done correctly and argue that increasing the number of sites is not expected to change our conclusions on the mass spectrum.

In the comment on our paper [1] Fujita, Kobayashi and Takahashi claim that our results for the mass spectrum of the massive Thirring model (MTM) are not reliable [2]. In particular, they claim that if one uses spin chain regularisation it is *necessary* to diagonalize the spin chain Hamiltonian with a number of sites N larger than 1000. We now explain why we disagree with this criticism.

The criticism of Fujita et al. is based on the following argument. If one makes the standard spin chain regularisation of MTM (which is XYZ spin $1/2$ chain [3, 4]) to obtain some reasonable results on continuum extrapolation, one has to satisfy the condition

$$\frac{2\pi}{N} \ll am_0 \ll 2\pi. \quad (1)$$

Here N is the number of sites, a is the lattice spacing, and m_0 is a bare mass parameter. Using (1) Fujita et al. claim that if one wants to obtain any reliable information on the bound state of the MTM one has to take $N > 1000$, which is much larger than ours $N \leq 16$. Moreover, they claim that for values of the parameters used in [1] the left inequality in (1) is even completely violated, i.e., $2\pi/N > am_0$. In addition they claim that the mass gap in our calculation is approximately equal to the “resolution” $2\pi/L$. From all this, Fujita et al. conclude that our results [1] cannot be very reliable for the bound state spectrum of the massive Thirring model.

Let us now review standard lattice philosophy. Continuum regime in lattice calculations is obtained when the correlation length ξ is much larger than the lattice spacing a and at the same time much smaller than spatial extension $L = Na$, i.e.,

$$a \ll \xi \ll Na. \quad (2)$$

As $\xi = 1/M$, where M is the mass gap (mass of the

lightest particle), (2) can be equivalently written as

$$\frac{1}{N} \ll Ma \ll 1 \quad (3)$$

Now, hardware limitations make restrictions on N . For example, in lattice (quenched) QCD maximum lattices which are presently calculable have $N \leq 64$ (see e.g., [5]). More importantly here, for exact diagonalisation in two dimensions $N < 30$. It follows that “ $\ll$ ” in (3) at best means 5–8 times smaller. So in practical calculations one effectively imposes the condition

$$\frac{1}{N} < Ma < 1 \quad (4)$$

and from the quality of scaling and the accuracy of the continuum extrapolation (usually using different methods as a check) one decides whether the extrapolated results are a good approximation of the continuum theory. Of course, one cannot exclude the possibility that for values of scaling parameter larger than accessed there is a complete change of scaling behavior so that the obtained extrapolated results are wrong[15].

Let us now apply the above textbook analysis to our lattice calculation (XYZ spin chain regularisation of MTM) [1]. For the sake of clarity we restrict ourselves to a interval of coupling constant where elementary fermion is the particle with the lowest mass. First, it is easy to see that (3) is *not* equal to (1) used by Fujita et al. It was shown in [4] that (using notation from [1])

$$m_0 a = \frac{8\gamma}{\pi} \sin \gamma \left(\frac{Ma}{4} \right)^{2\gamma/\pi}.$$

In particular, let us analyze the left inequality in (4). In our extrapolation we had $Ma > 0.2$ which is for $N = 16$ reasonably larger than $1/N = 0.06$. It follows that in our analysis continuum condition is fairly satisfied when *proper* condition is used, contrary to claim of Fujita et al.

As we mentioned above, there is always the possibility that for larger N something dramatic happens with the scaling law and that our extrapolations made with $N \leq$

*Electronic address: mk679@nyu.edu

†Electronic address: pallua@phy.hr

‡Electronic address: pprester@phy.hr

16 are incorrect. There are several reasons why we believe that this should not be expected in the case of the XYZ chain:

- global properties of the energy spectrum are as expected for QFT with calculated mass spectrum. In particular, there are states with energies corresponding to two-fermion and two-(first)breather states
- the same technique was previously successfully applied for similar perturbed CFT's, e.g. ordinary and tricritical Ising model in a magnetic field [7]. It was shown there that passing from $N \leq 14$ to $N \leq 21$ only slightly improved precision and conclusions about the spectra remained the same. In fact it was shown that very accurate results can be obtained already for lattices with $N \leq 24$ [8].
- our results agree with the DHN mass formula [9] obtained by a number of different methods. It is very hard to imagine how bad extrapolation can agree with exact analytic result. We should also mention that a similar analysis for lattice regularization of the sine-Gordon model (periodic XXZ chain in transverse magnetic field) also gave results consistent with the DHN formula [10].
- Beside the mass ratios in $L \rightarrow \infty$ limit we obtained anomalous dimensions of corresponding states, in particular we had for the first time calculated dimension of the second breather. Our result was subsequently confirmed by analytical calculation [11].

We conclude that our numerical analysis made in [1] was done correctly and that all results should be trusted,

in a sense that enlarging lattice would just increase numerical precision and leave all conclusions unchanged.

Finally, we would like to comment on the Bethe ansatz solution of the MTM. Spectrum of the MTM can be found by solving the Bethe ansatz equations in the continuum approximation [12]. The spectrum found there agrees with the results of [9]. Recently, it was argued [13] that there exist no complex solution to the Bethe equations and that there is only one bound state. In [12] precisely the complex solutions (so called Bethe strings) are describing bound states of MTM. We performed numerical analysis of Bethe equations for MTM and tried to reproduce the result of [13]. Our results indicate that the powers α that determine the scaling behavior of energies of the few lowest states on the density ρ differ and even vary with the ρ (for definitions see section 4 in [13]). Therefore we were unable to extract conclusive results by letting $\rho \rightarrow \infty$. An indication that the numerical iterations might easily miss the complex solutions (that nevertheless exist) is coming from our study of the Bethe ansatz equations for different spin chains [14]. There we studied how the complex (string) solution in two and three particle sectors emerge and disappear when one changes the parameters of the model. We found that iterative numerical methods fail to converge on string solutions, although they exist and can be found analytically.

Acknowledgments

MK would like to thank to J.H. Lowenstein for useful and stimulating discussion.

-
- [1] M. Kolanović, S. Pallua and P. Prester, Phys. Rev. D **62**, 025021 (2000).
 - [2] T. Fujita, T. Kobayashi and H. Takahashi, hep-th/0306173.
 - [3] A. Luther, Phys. Rev. B **14**, 2153 (1976).
 - [4] M. Lüscher, Nucl. Phys. **B117**, 475 (1976).
 - [5] S. Aoki et al. (CP-PACS Collab.), hep-lat/0206009.
 - [6] P. Christe and M. Henkel, *Introduction to Conformal Invariance and Its Applications to Critical Phenomena* (Springer-Verlag, Heidelberg, 1993).
 - [7] M. Henkel and H. Saleur, J. Phys. A **22**, L513 (1989); M. Henkel, J. Phys. A **24**, L133 (1991); M. Henkel, J. Phys. A **23**, 4369 (1990).
 - [8] M. Caselle, M. Hasenbusch, A. Pelissetto and E. Vicari, J. Phys. A **33**, 8171 (2000); M. Caselle and M. Hasenbusch, Nucl. Phys. **B579**, 667 (2000).
 - [9] R. F. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. D **11**, 3424 (1975).
 - [10] S. Pallua and P. Prester, Phys. Rev. D **59**, 125006 (1999); S. Pallua and P. Prester, Fiz. **B10**, 175 (2001).
 - [11] G. Feverati, F. Ravanini, and G. Takacs, Nucl. Phys. **B570**, 615 (2000).
 - [12] H. Bergknoff and H. B. Thacker, Phys. Rev. D **19**, 3666 (1979); H. Bergknoff and H. B. Thacker, Phys. Rev. Lett. **42**, 135 (1979).
 - [13] T. Fujita, Y. Sekiguchi and K. Yamamoto, Annals Phys. **255**, 204 (1997).
 - [14] A. Ilakovac, M. Kolanović, S. Pallua and P. Prester, Phys. Rev. B **60**, 7271 (1999); A. Ilakovac, M. Kolanović, S. Pallua and P. Prester, Fiz. **B8**, 453 (1999).
 - [15] For nice discussion see Sec. 9.5 in Ref. [6].